

## **Acknowledgments**

The report is the summarization of many projects. Our projects are sponsored by Ministry of Science and Technology, State Administration of Forestry, Anhui Province, Guangxi Zhuang Autonomous Region, Guangdong Province, Zhejiang Province and Liaoning Province and Committee on Trade and Investment (Forestry EVSL Export Group), APEC. APEC finances us to make this publication. Many colleagues, friends and associates have contributed to this brochure by providing data, methods and ideas. We wish to express our appreciation to all people who have helped and all units that have supported our work.

## References

- Brockhaus J A.£-Khorram S. et al (1992). A comparison of Landsat TM and SPOT HRV data for use in the development of forest defoliation modes. *International Journal of Remote Sensing*, 13(6):3235-3240.
- Ekstrand S P. (1990). Detection of Moderate Damage on Norway Spruce Using Landsat TM and Digital Stand Data. *IEEE Transactions on Geoscience and Remote Sensing*, 28(4):685- 692.
- Gills M D., Pick R D. and Leckie D G. (1990). Satellite imagery assists in the assessment of hail damage for salvage harvest. *The Forestry Chronicle*, (10):463-468.
- Joria P E., Ahearn S C. et al. (1991). A Comparison of the SPOT and Landsat Thematic Mapper Satellite Systems for Detecting Gypsy Moth Defoliation in Michigan. *Photogrammetric Engineering & Remote Sensing*, 57(12):1605-1612
- Leckie Donald G, Yuan Xiaoping et al. (1992). Analysis of High Resolution Multispectral MEIS Imagery for Spruce Budworm Damage Assessment on a Single Tree Basis. *Remote Sensing of Environment*, 40:125-136
- Linden D S., Hoffer R M., pywell H R. (1996). Automated Digital Mosaicking of Airborne Videography. Lothar, B., *Satellite remote sensing forest atlas of Europe*. 1995
- McNairn H., Protz H. (1993). Mapping corn residue cover on agricultural fields in Oxford county, Ontario, using Thematic Mapper. *Canadian Journal of Remote Sensing*, 19 (2):152 -159
- Mehrotra,A. and Suri, R.K., *Remote sensing for environment and forest management*. 1994
- Nakane K, Kimura Y.(1992). Assessment of pine forest damage by blight based on Landsat TM data and correlation with environmental factors. *Ecological Research*, (7) :9-18
- Prakasa B.L.S. (1993) *Nonparametric Functional Estimation*, Academic Press Inc.
- Rock, B.N., Vogelmann, J.E., Williams, D.L., Vogelmann, A.F. and Hoshizaki, T., 1986. Remote sensing of forest damage, *Bioscience*. 36(7):439-445
- Vogelmann J E, Rock B N.(1986). Assessing forest decline in coniferous of Vermont using ZS-001 Thematic Mapper Simulator data. *International Journal of Remote Sensing*, 7(10): 1303 -1321
- Vogelmann J E, Rock B N.(1988). Assessing forest damage in high-elevation coniferous forests in Vermont and New Hampshire using Thematic Mapper data. *Remote Sensing of Environment*, 24:227-246
- Vogelmann J E.(1990). Comparison between two vegetation indices for measuring different types of forest damage in the north-eastern United States. *International Journal of Remote Sensing*, 11(12):2281-2297
- Vogelmann J E., Rock B N. et al.(1989). Use of Thematic Mapper Data for the Detection of Forest Damage Caused by the Pear Thrips. *Remote Sensing of Environment*, 30: 217-225
- Vogelmann J.E. ,1990, Comparison between two vegetation indices for measuring different types of forest damage in the north-eastern United States, *International Journal of Remote Sensing*, 11(12):2281-2297
- William, M.C., *Remote sensing in forest health protection*. 2000
- Zhang X Y., Lu Q. AND Chao L J. The research of the science of forest protection towards the 21st century workshop on forestry and agroforestry for environmental protection and rural development 2001